

Trigonometric Identities

Jonathan Senning, jonathan.senning@gordon.edu

September 1999. Revised 2000, 2003, 2006, 2010.

1 Trigonometric Identities you must remember

The “big three” trigonometric identities are

$$\sin^2 \theta + \cos^2 \theta = 1 \tag{1}$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta \tag{2}$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \tag{3}$$

Using these we can derive many other identities. Even if we commit the other useful identities to memory, these three will help be sure that our signs are correct, etc.

2 Two more easy identities

From equation (1) we can generate two more identities. First, divide each term in (1) by $\cos^2 \theta$ (assuming it is not zero) to obtain

$$\tan^2 \theta + 1 = \sec^2 \theta. \tag{4}$$

When we divide by $\sin^2 \theta$ (again assuming it is not zero) we get

$$1 + \cot^2 \theta = \csc^2 \theta. \tag{5}$$

3 Identities involving the difference of two angles

From equations (2) and (3) we can get several useful identities. First, recall that

$$\cos(-\theta) = \cos \theta, \quad \sin(-\theta) = -\sin \theta. \tag{6}$$

From (2) we see that

$$\begin{aligned} \sin(\alpha - \beta) &= \sin(\alpha + (-\beta)) \\ &= \sin \alpha \cos(-\beta) + \cos \alpha \sin(-\beta) \end{aligned}$$

which, using the relationships in (6), reduces to

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta. \tag{7}$$

In a similar way, we can use equation (3) to find

$$\begin{aligned} \cos(\alpha - \beta) &= \cos(\alpha + (-\beta)) \\ &= \cos \alpha \cos(-\beta) - \sin \alpha \sin(-\beta) \end{aligned}$$

which simplifies to

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta. \tag{8}$$

Notice that by remembering the identities (2) and (3) you can easily work out the signs in these last two identities.

4 Identities involving products of sines and cosines

If we now add equation (2) to equation (7)

$$\begin{aligned}\sin(\alpha - \beta) &= \sin \alpha \cos \beta - \cos \alpha \sin \beta \\ +(\sin(\alpha + \beta) &= \sin \alpha \cos \beta + \cos \alpha \sin \beta)\end{aligned}$$

we find

$$\sin(\alpha - \beta) + \sin(\alpha + \beta) = 2 \sin \alpha \cos \beta$$

and dividing both sides by 2 we obtain the identity

$$\sin \alpha \cos \beta = \frac{1}{2} \sin(\alpha - \beta) + \frac{1}{2} \sin(\alpha + \beta). \quad (9)$$

In the same way we can add equations (3) and (8)

$$\begin{aligned}\cos(\alpha - \beta) &= \cos \alpha \cos \beta + \sin \alpha \sin \beta \\ +(\cos(\alpha + \beta) &= \cos \alpha \cos \beta - \sin \alpha \sin \beta)\end{aligned}$$

to get

$$\cos(\alpha - \beta) + \cos(\alpha + \beta) = 2 \cos \alpha \cos \beta$$

which can be rearranged to yield the identity

$$\cos \alpha \cos \beta = \frac{1}{2} \cos(\alpha - \beta) + \frac{1}{2} \cos(\alpha + \beta). \quad (10)$$

Suppose we wanted an identity involving $\sin \alpha \sin \beta$. We can find one by slightly modifying the last thing we did. Rather than adding equations (3) and (8), all we need to do is subtract equation (3) from equation (8):

$$\begin{aligned}\cos(\alpha - \beta) &= \cos \alpha \cos \beta + \sin \alpha \sin \beta \\ -(\cos(\alpha + \beta) &= \cos \alpha \cos \beta - \sin \alpha \sin \beta)\end{aligned}$$

This gives

$$\cos(\alpha - \beta) - \cos(\alpha + \beta) = 2 \sin \alpha \sin \beta$$

or, in the form we prefer,

$$\sin \alpha \sin \beta = \frac{1}{2} \cos(\alpha - \beta) - \frac{1}{2} \cos(\alpha + \beta). \quad (11)$$

5 Double angle identities

Now a couple of easy ones. If we let $\alpha = \beta$ in equations (2) and (3) we get the two identities

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha, \quad (12)$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha. \quad (13)$$

6 Identities for sine squared and cosine squared

If we have $\alpha = \beta$ in equation (10) then we find

$$\begin{aligned}\cos \alpha \cos \beta &= \frac{1}{2} \cos(\alpha - \alpha) + \frac{1}{2} \cos(\alpha + \alpha) \\ \cos^2 \alpha &= \frac{1}{2} \cos 0 + \frac{1}{2} \cos 2\alpha.\end{aligned}$$

Simplifying this and doing the same with equation (11) we find the two identities

$$\cos^2 \alpha = \frac{1}{2}(1 + \cos 2\alpha), \quad (14)$$

$$\sin^2 \alpha = \frac{1}{2}(1 - \cos 2\alpha). \quad (15)$$

7 Identities involving tangent

Finally, from equations (2) and (3) we can obtain an identity for $\tan(\alpha + \beta)$:

$$\tan(\alpha + \beta) = \frac{\sin(\alpha + \beta)}{\cos(\alpha + \beta)} = \frac{\sin \alpha \cos \beta + \cos \alpha \sin \beta}{\cos \alpha \cos \beta - \sin \alpha \sin \beta}.$$

Now divide numerator and denominator by $\cos \alpha \cos \beta$ to obtain the identity we wanted:

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}. \quad (16)$$

We can get the identity for $\tan(\alpha - \beta)$ by replacing β in (16) by $-\beta$ and noting that tangent is an odd function:

$$\tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}. \quad (17)$$

8 Summary

There are many other identities that can be generated this way. In fact, the derivations above are not unique — many trigonometric identities can be obtained many different ways. The idea here is to be very familiar with a small number of identities so that you are comfortable manipulating and combining them to obtain whatever identity you need to.